

11.1. Let G be a locally compact group. As was shown in the lectures, $L^1(G)$ is a Banach algebra under convolution.

(a) Show that $L^1(G)$ is a Banach $*$ -algebra w.r.t. the involution $f^*(x) = \overline{f(x^{-1})}\Delta(x^{-1})$ ($f \in L^1(G)$, $x \in G$).

(b) Show that $L^1(G)$ (equipped with the standard L^1 -norm and with the involution defined in (a)) is not a C^* -algebra unless $G = \{e\}$.

(c) Show that $L^1(G)$ is commutative if and only if G is commutative.

(d) Show that $L^1(G)$ is unital if and only if G is discrete.

11.2. Let G be a locally compact group, and let $p, q \in (1, +\infty)$ satisfy $1/p + 1/q = 1$. Show that, for each $f \in L^p(G)$ and $g \in L^q(G)$, the convolution $f * Sg$ (where $(Sg)(x) = g(x^{-1})$) is defined everywhere on G , belongs to $C_0(G)$, and that $\|f * Sg\|_\infty \leq \|f\|_p \|g\|_q$.

11.3. Let G be a locally compact group.

(a) Show that $M(G)$ is a unital Banach $*$ -algebra w.r.t. the involution $\nu^*(B) = \overline{\nu(B^{-1})}$ ($\nu \in M(G)$, $B \subset G$ is a Borel set). In particular, show that convolution is associative on $M(G)$ (this was not proved at the lectures).

(b) Show that $M(G)$ is commutative if and only if G is commutative.

(c) Let μ be a left Haar measure on G . Show that the map $i: L^1(G) \rightarrow M(G)$, $f \mapsto f \cdot \mu$, is an isometric $*$ -algebra homomorphism.

(d) Identify $L^1(G)$ with its canonical image in $M(G)$ (see (c) above). Show that $L^1(G)$ is a closed 2-sided ideal of $M(G)$, and that for each $f \in L^1(G)$, $\nu \in M(G)$, and for almost all $x \in G$ we have

$$(\nu * f)(x) = \int_G f(y^{-1}x) d\nu(y), \quad (f * \nu)(x) = \int_G f(xy^{-1})\Delta(y^{-1}) d\nu(y). \quad (1)$$

11.4. Let G be a locally compact group, and let λ (resp. ρ) denote the left (resp. right) regular representation of G on $L^1(G)$. Show that $\lambda(x)f = \delta_x * f$ and $\rho(x)f = f * \delta_{x^{-1}}$ ($f \in L^1(G)$, $x \in G$).

11.5. Let X be a locally compact Hausdorff topological space.

(a) Construct a bounded approximate identity in $C_0(X)$.

(b) Show that $C_0(X)$ has a sequential bounded approximate identity if and only if X is σ -compact.

11.6. Let H be a Hilbert space.

(a) Construct a bounded approximate identity in $\mathcal{K}(H)$.

(b) Show that $\mathcal{K}(H)$ has a sequential bounded approximate identity if and only if H is separable.

11.7. Let G be a locally compact group, and let (u_i) be a Dirac net in $L^1(G)$. Show that (u_i) converges to $\delta_e \in M(G)$ w.r.t. the weak* topology on $M(G)$.

11.8. Let A be a Banach algebra with a bounded approximate identity (e_α) , and let E be a left Banach A -module. Recall that the *essential part* of E is $E_{\text{ess}} = \overline{\text{span}\{av : a \in A, v \in E\}}$.

(a) Show that E_{ess} is the largest essential submodule of E .

(b) Show that $E_{\text{ess}} = \{v \in E : v = \lim e_\alpha v\}$.

(c) Does (a) hold without the assumption that A has a b.a.i.?

11.9. Let G be a locally compact group, $\nu \in M(G)$, and $f \in L^p(G)$ (where $1 \leq p \leq \infty$).

(a) Show that the convolution $\nu * f$ given by (1) is defined a.e. on G , that $\nu * f \in L^p(G)$, and that the action $(\nu, f) \mapsto \nu * f$ makes $L^p(G)$ into a left unital Banach $M(G)$ -module.

(b) Show that $L^p(G)$ is essential over $L^1(G)$ if $p < \infty$.

(c) Find the essential part of $L^\infty(G)$ over $L^1(G)$.

11.10. Define $f: [0, 1] \rightarrow c_0$ by

$$f(t) = (\chi_{(0,1]}(t), 2\chi_{(0,1/2]}(t), \dots, n\chi_{(0,1/n]}(t), \dots) \quad (t \in [0, 1]).$$

Show that f is Dunford integrable (w.r.t. the Lebesgue measure on $[0, 1]$), but is not Pettis integrable.

11.11. Let (X, μ) be a measure space, and let E, F be Banach spaces.

(a) Suppose that $f: X \rightarrow E$ is Dunford integrable and that $\|f\|: x \mapsto \|f(x)\|$ is integrable. Show that $\|\int f d\mu\| \leq \int \|f\| d|\mu|$.

(b) Suppose that $f: X \rightarrow E$ is Pettis integrable. Show that for each bounded linear map $T: E \rightarrow F$ the function $T \circ f$ is Pettis integrable, and that $T(\int f d\mu) = \int (T \circ f) d\mu$.

11.12. Let G be a locally compact group, and let π be a uniformly bounded continuous representation of G on a reflexive Banach space E . Recall that the *canonical extension* of π to $M(G)$ is given by $\tilde{\pi}(\nu)v = \int_G \pi(x)v d\nu(x)$ ($\nu \in M(G)$, $v \in E$).

(a) Show that $\tilde{\pi}$ is indeed a representation of $M(G)$ on E .

(b) Choose a left Haar measure μ on G . Show that for each $g \in L^1(G)$ (where $L^1(G)$ is canonically embedded into $M(G)$) we have $\tilde{\pi}(g)v = \int_G g(x)\pi(x)v d\mu(x)$.

(c) Suppose that E is a Hilbert space. Show that $\tilde{\pi}$ is a $*$ -representation if and only if π is unitary.

11.13. Let G be a locally compact group. Show that, for each $\nu \in M(G)$ and $f \in L^p(G)$ (where $1 < p < \infty$), we have $\nu * f = \tilde{\lambda}(\nu)f$, where λ is the left regular representation of G on $L^p(G)$ and $\tilde{\lambda}$ is the canonical extension of λ to $M(G)$.

11.14. Let A be a Banach algebra, and let $B \subset A$ be a closed 2-sided ideal with a bounded approximate identity. Show that

(a) if E is a B -essential Banach A -module and $E_0 \subset E$ is a closed B -submodule, then E_0 is an A -submodule;

(b) if E and F are B -essential Banach A -modules, then $\text{Hom}_A(E, F) = \text{Hom}_B(E, F)$;

(c) if H is a Hilbert space equipped with an action of A which makes H into a B -essential Banach A -module, then H is a $*$ -module over B iff H is a $*$ -module over A .

11.15. Define a representation π of \mathbb{R} on $L^2(\mathbb{R})$ by $(\pi(t)f)(x) = e^{-2\pi itx}f(x)$. Find an explicit formula for the associated representation $\tilde{\pi}$ of $L^1(\mathbb{R})$. Show that π is unitarily isomorphic to the left regular representation.