

3.1. As in Exercises 2.8 and 2.9, define the convolution product on $L^1(\mathbb{R})$, show that $L^1(\mathbb{R})$ is a commutative nonunital algebra, and prove that the Fourier transform $\mathcal{F}_{\mathbb{R}}: L^1(\mathbb{R}) \rightarrow C_0(\mathbb{R})$ is an algebra homomorphism.

3.2. Suppose that $f \in C^1(\mathbb{R})$ and that $f, f' \in L^1(\mathbb{R})$. Prove that $(f')^\wedge(\lambda) = 2\pi i \lambda \hat{f}(\lambda)$ ($\lambda \in \mathbb{R}$). Deduce that if $f \in C^p(\mathbb{R})$ and $f, f', \dots, f^{(p)} \in L^1(\mathbb{R})$, then $\hat{f}(\lambda) = o(|\lambda|^{-p})$ as $\lambda \rightarrow \infty$.

3.3. Formulate and prove a result similar to Exercise 3.2 for the Fourier transform on \mathbb{T} .

3.4. Let $t = \mathbf{1}_{\mathbb{R}}$ denote the identity map on \mathbb{R} . Let $f \in L^1(\mathbb{R})$, and suppose that $tf \in L^1(\mathbb{R})$. Show that $\hat{f} \in C^1(\mathbb{R})$, and that $\hat{f}'(\lambda) = -2\pi i (tf)^\wedge(\lambda)$ ($\lambda \in \mathbb{R}$). Deduce that if $f, tf, \dots, t^p f \in L^1(\mathbb{R})$, then $\hat{f} \in C^p(\mathbb{R})$.

3.5. Formulate and prove a result similar to Exercise 3.4 for the Fourier transform on \mathbb{Z} .

3.6. Let $\mathcal{F}: L^1(\mathbb{R}) \rightarrow C_0(\mathbb{R})$ denote the Fourier transform, and let $\hat{\mathcal{F}} = S\mathcal{F}$, where $(Sf)(t) = f(-t)$ ($t \in \mathbb{R}$).

(a) Show that \mathcal{F} and $\hat{\mathcal{F}}$ map the Schwartz space $\mathcal{S}(\mathbb{R})$ continuously into itself.

(b) Suppose that $T: \mathcal{S}(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{R})$ is a linear map commuting with $\frac{d}{dt}$ and with the multiplication by the coordinate t . Show that $T = c\mathbf{1}_{\mathcal{S}(\mathbb{R})}$ for some $c \in \mathbb{C}$.

(c) Let $f(t) = e^{-\pi t^2}$ ($t \in \mathbb{R}$). Show that $\hat{f} = f$.

(d) Deduce from (a), (b), (c) that $\mathcal{F}\hat{\mathcal{F}} = \hat{\mathcal{F}}\mathcal{F} = \mathbf{1}_{\mathcal{S}(\mathbb{R})}$ on $\mathcal{S}(\mathbb{R})$. In other words, \mathcal{F} is a topological isomorphism of $\mathcal{S}(\mathbb{R})$ onto itself, and $\mathcal{F}^2 = S$ on $\mathcal{S}(\mathbb{R})$.

3.7. (This is an analog of Exercise 3.6 for \mathbb{Z} and \mathbb{T} .) Let $C_{2\pi}^\infty(\mathbb{R})$ denote the space of all smooth 2π -periodic functions on \mathbb{R} , and let $j: C^\infty(\mathbb{T}) \rightarrow C_{2\pi}^\infty(\mathbb{R})$ denote the vector space isomorphism given by $(jf)(t) = f(e^{it})$ ($t \in \mathbb{R}$). Given $f \in C^\infty(\mathbb{T})$, define the derivative $f' \in C^\infty(\mathbb{T})$ of f by $f' = j^{-1}(j(f)')$. The higher derivatives $f^{(k)}$ are defined in an obvious way. We endow $C^\infty(\mathbb{T})$ with the topology generated by the family $\{\|\cdot\|_k : k \in \mathbb{Z}_{\geq 0}\}$ of seminorms, where $\|f\|_k = \sup_{z \in \mathbb{T}} |f^{(k)}(z)|$.

We define the space of *rapidly decreasing sequences* by

$$s(\mathbb{Z}) = \left\{ x = (x_n) \in \mathbb{C}^{\mathbb{Z}} : \|x\|_k = \sup_{n \in \mathbb{Z}} |x_n| |n|^k < \infty \forall k \in \mathbb{Z}_{\geq 0} \right\}$$

and topologize $s(\mathbb{Z})$ by the family $\{\|\cdot\|_k : k \in \mathbb{Z}_{\geq 0}\}$ of seminorms. Prove that

(a) $\mathcal{F}_{\mathbb{Z}}$ maps $s(\mathbb{Z})$ continuously into $C^\infty(\mathbb{T})$;

(b) $\mathcal{F}_{\mathbb{T}}$ maps $C^\infty(\mathbb{T})$ continuously into $s(\mathbb{Z})$;

(c) $\mathcal{F}_{\mathbb{T}}\mathcal{F}_{\mathbb{Z}} = S_{\mathbb{Z}}$ and $\mathcal{F}_{\mathbb{Z}}\mathcal{F}_{\mathbb{T}} = S_{\mathbb{T}}$, where $(S_{\mathbb{Z}}f)(n) = f(-n)$ and $(S_{\mathbb{T}}g)(z) = g(z^{-1})$ for every $f \in s(\mathbb{Z})$ and $g \in C^\infty(\mathbb{T})$. As a consequence, $\mathcal{F}_{\mathbb{Z}}$ and $\mathcal{F}_{\mathbb{T}}$ are topological isomorphisms between $s(\mathbb{Z})$ and $C^\infty(\mathbb{T})$.

3.8. Given $\lambda \in \mathbb{R}$, let $\chi_\lambda(t) = e^{-2\pi i \lambda t}$ ($t \in \mathbb{R}$). (Recall that the χ_λ 's are precisely the unitary characters of \mathbb{R} .) Find the Fourier transforms of χ_λ and of the Dirac δ -function δ_λ .

3.9. Let $s'(\mathbb{Z})$ denote the topological dual of $s(\mathbb{Z})$ (i.e., the space of all continuous linear functionals on $s(\mathbb{Z})$). Show that the map $\varphi \mapsto (\varphi(\delta_n))_{n \in \mathbb{Z}}$ is a vector space isomorphism between $s'(\mathbb{Z})$ and the space of *tempered sequences*

$$\left\{ x = (x_n) \in \mathbb{C}^{\mathbb{Z}} : |x_n| |n|^{-k} \text{ is bounded for some } k \in \mathbb{Z}_{\geq 0} \right\}.$$

3.10. Let $\mathcal{D}'(\mathbb{T})$ denote the topological dual of $C^\infty(\mathbb{T})$ (i.e., the space of all continuous linear functionals on $C^\infty(\mathbb{T})$). The elements of $\mathcal{D}'(\mathbb{T})$ are called *distributions* on \mathbb{T} . Given $f \in L^1(\mathbb{T})$, define $\varphi_f \in \mathcal{D}'(\mathbb{T})$ by $\varphi_f(g) = \int_{\mathbb{T}} fg d\mu$. Show that the map $L^1(\mathbb{T}) \rightarrow \mathcal{D}'(\mathbb{T})$, $f \mapsto \varphi_f$, is injective.

3.11. Define the Fourier transforms $\mathcal{F}_{\mathbb{Z}}: s'(\mathbb{Z}) \rightarrow \mathcal{D}'(\mathbb{T})$ and $\mathcal{F}_{\mathbb{T}}: \mathcal{D}'(\mathbb{T}) \rightarrow s'(\mathbb{Z})$ to be the maps dual to $\mathcal{F}_{\mathbb{T}}: C^\infty(\mathbb{T}) \rightarrow s(\mathbb{Z})$ and $\mathcal{F}_{\mathbb{Z}}: s(\mathbb{Z}) \rightarrow C^\infty(\mathbb{T})$, respectively.

(a) Identify $c_0(\mathbb{Z})$ with a subspace of $s'(\mathbb{Z})$ via Exercise 3.9, and identify $L^1(\mathbb{T})$ with a subspace of $\mathcal{D}'(\mathbb{T})$ via Exercise 3.10. Show that that the Fourier transforms on $s'(\mathbb{Z})$ and on $\mathcal{D}'(\mathbb{T})$ extend the “classical” Fourier transforms $\ell^1(\mathbb{Z}) \rightarrow C(\mathbb{T})$ and $L^1(\mathbb{T}) \rightarrow c_0(\mathbb{Z})$.

(b) (This is an analog of Exercise 3.8.) Calculate the Fourier transforms of the unitary characters and of the Dirac δ -functions on \mathbb{Z} and on \mathbb{T} .

(c) (*the Fourier series in $\mathcal{D}'(\mathbb{T})$*). Show that for each $f \in \mathcal{D}'(\mathbb{T})$ we have $f = \sum_{n \in \mathbb{Z}} \hat{f}(n)\chi_{-n}$, where the series converges in the weak* topology on $\mathcal{D}'(\mathbb{T})$ (i.e., the topology of pointwise convergence on elements of $C^\infty(\mathbb{T})$).

3.12. (a) Define a canonical topology on $C^\infty(\mathbb{T}^2)$ by analogy with $C^\infty(\mathbb{T})$.

(b) Show that the map

$$C^\infty(\mathbb{T}) \otimes C^\infty(\mathbb{T}) \rightarrow C^\infty(\mathbb{T}^2), \quad f \otimes g \mapsto ((z, w) \mapsto f(z)g(w)),$$

is injective and has dense image. From now on, we identify $C^\infty(\mathbb{T}) \otimes C^\infty(\mathbb{T})$ with a dense subspace of $C^\infty(\mathbb{T}^2)$ via the above map.

(c) (*tensor product of distributions*). For each φ, ψ in $\mathcal{D}'(\mathbb{T})$ the element $\varphi \otimes \psi \in \mathcal{D}'(\mathbb{T}) \otimes \mathcal{D}'(\mathbb{T})$ may be viewed as a linear functional on $C^\infty(\mathbb{T}) \otimes C^\infty(\mathbb{T})$. Show that $\varphi \otimes \psi$ uniquely extends to a continuous linear functional on $C^\infty(\mathbb{T}^2)$.

(d) Define $\Delta: C^\infty(\mathbb{T}) \rightarrow C^\infty(\mathbb{T}^2)$ by $(\Delta f)(z, w) = f(zw)$. For each φ, ψ in $\mathcal{D}'(\mathbb{T})$ define the *convolution* $\varphi * \psi \in \mathcal{D}'(\mathbb{T})$ by

$$\langle \varphi * \psi, f \rangle = \langle \varphi \otimes \psi, \Delta f \rangle \quad (f \in C^\infty(\mathbb{T})).$$

Show that $(\mathcal{D}'(\mathbb{T}), *)$ is a commutative unital algebra containing $L^1(\mathbb{T})$ and $\mathbb{C}\mathbb{T}$ as subalgebras. In particular, the convolution on $\mathcal{D}'(\mathbb{T})$ agrees with those on $L^1(\mathbb{T})$ and on $\mathbb{C}\mathbb{T}$.

(e) Identify $s'(\mathbb{Z})$ with the space of tempered sequences (see Exercise 3.9). Show that $s'(\mathbb{Z})$ is a unital algebra under pointwise multiplication, and that the Fourier transforms $\mathcal{F}_{\mathbb{Z}}$ and $\mathcal{F}_{\mathbb{T}}$ (see Exercise 3.11) are algebra isomorphisms between $s'(\mathbb{Z})$ and $\mathcal{D}'(\mathbb{T})$.

3.13 (*the Poisson summation formula*). Identify \mathbb{T} with \mathbb{R}/\mathbb{Z} , and define $a: \mathcal{S}(\mathbb{R}) \rightarrow C^\infty(\mathbb{T})$ by $(af)(t + \mathbb{Z}) = \sum_{n \in \mathbb{Z}} f(t + n)$. Show that we indeed have $af \in C^\infty(\mathbb{T})$ whenever $f \in \mathcal{S}(\mathbb{R})$, and that the diagram

$$\begin{array}{ccc} \mathcal{S}(\mathbb{R}) & \xrightarrow{\mathcal{F}_{\mathbb{R}}} & \mathcal{S}(\mathbb{R}) \\ a \downarrow & & \downarrow \text{restr.} \\ C^\infty(\mathbb{T}) & \xrightarrow{\mathcal{F}_{\mathbb{T}}} & s(\mathbb{Z}) \end{array}$$

commutes. Deduce that for each $f \in \mathcal{S}(\mathbb{R})$ we have

$$\sum_{n \in \mathbb{Z}} f(t + n) = \sum_{n \in \mathbb{Z}} \hat{f}(n)e^{2\pi int} \quad (t \in \mathbb{R}).$$