

## Topological vector spaces

Exercises marked by “-B” are optional. If you solve such exercises, you will earn bonus points.

**3.1.** Let  $X$  and  $Y$  be topological vector spaces. Show that

- (a) a linear operator  $X \rightarrow Y$  is continuous iff it is continuous at 0;  
 (b) the set  $\mathcal{L}(X, Y)$  of continuous linear operators from  $X$  to  $Y$  is a vector subspace of the space of all linear operators from  $X$  to  $Y$ .

**3.2.** Is there at least one continuous norm on the following topological vector spaces?

- (a)  $\mathbb{K}^X$  (where  $X$  is a set); (b)  $C(X)$  (where  $X$  is a metrizable topological space);  
 (c) the space  $\mathcal{O}(U)$  of holomorphic functions on an open set  $U \subset \mathbb{C}$  (we equip  $\mathcal{O}(U)$  with the topology induced from  $C(U)$ );  
 (d)  $C^\infty[a, b]$ ; (e)  $C^\infty(U)$ , where  $U \subset \mathbb{R}^n$  is an open set; (f)  $\mathcal{S}(\mathbb{R}^n)$ .

**3.3.** Let  $X$  be a Hausdorff locally convex space, and let  $P$  be a defining family of seminorms on  $X$ . Show that  $X$  is normable iff  $P$  is equivalent to a finite subfamily  $P_0 \subset P$ .

**3.4. (a)-(f)** Which spaces of Exercise 3.2 are normable?

**3.5.** Let  $X$  be a Hausdorff locally convex space, and let  $P$  be a defining family of seminorms on  $X$ . Show that  $X$  is metrizable iff  $P$  is equivalent to an at most countable subfamily  $P_0 \subset P$ .

*Hint.* If  $(p_n)_{n \in \mathbb{N}}$  is a sequence of seminorms, then the function

$$\rho(x, y) = \sum_n \frac{1}{2^n} \min\{p_n(x - y), 1\} \quad \text{or, if you like,} \quad \rho(x, y) = \sum_n \frac{1}{2^n} \frac{p_n(x - y)}{1 + p_n(x - y)}$$

satisfies the triangle inequality.

**3.6. (a)-(f)** Which spaces of Exercise 3.2 are metrizable?

**3.7-B.** Let  $X$  be a finite-dimensional vector space. Show that there is only one topology on  $X$  making  $X$  into a Hausdorff locally convex space.

**3.8.** Let  $X$  be a set. Prove that for each  $f \in \mathbb{K}^X$  the multiplication operator  $M_f: \mathbb{K}^X \rightarrow \mathbb{K}^X$ ,  $M_f(g) = fg$ , is continuous.

**3.9.** Let  $U \subset \mathbb{R}^n$  be an open set. Prove that each linear differential operator  $\sum_{|\alpha| \leq N} a_\alpha D^\alpha$  on  $C^\infty(U)$  (where  $a_\alpha \in C^\infty(U)$ ) is continuous.

**3.10.** Let  $\mathbb{D}_R = \{z \in \mathbb{C} : |z| < R\}$ . Given  $f \in \mathcal{O}(\mathbb{D}_R)$ , let  $c_n(f) = f^{(n)}(0)/n!$ . Show that the topology of compact convergence on  $\mathcal{O}(\mathbb{D}_R)$  is generated by the family  $\{\|\cdot\|_{r, \infty} : 0 < r < R\}$  of seminorms, where  $\|f\|_{r, \infty} = \sup_{n \geq 0} |c_n(f)|r^n$ .

**3.11.** Recall (see the lectures) that the standard topology on the Schwartz space  $\mathcal{S}(\mathbb{R})$  is given by the family  $\{\|\cdot\|_{k, \ell} : k, \ell \in \mathbb{Z}_{\geq 0}\}$  of seminorms, where  $\|\varphi\|_{k, \ell} = \sup_{x \in \mathbb{R}} |x^k \varphi^{(\ell)}(x)|$ . Show that the following families of seminorms generate the same topology on  $\mathcal{S}(\mathbb{R})$ :

- (a)  $\{\|\cdot\|_\infty^{(p)} : p \in \mathbb{Z}_{\geq 0}\}$ , where  $\|\varphi\|_\infty^{(p)} = \sup_{k \leq p, x \in \mathbb{R}} (1 + x^2)^{p/2} |\varphi^{(k)}(x)|$ ;  
 (b)  $\{\|\cdot\|_1^{(p)} : p \in \mathbb{Z}_{\geq 0}\}$ , where  $\|\varphi\|_1^{(p)} = \max_{k \leq p} \int_{\mathbb{R}} (1 + x^2)^{p/2} |\varphi^{(k)}(x)| dx$ .

**3.12-B.** The space  $s(\mathbb{Z})$  of *rapidly decreasing sequences* is defined as follows:

$$s(\mathbb{Z}) = \left\{ x = (x_n) \in \mathbb{K}^{\mathbb{Z}} : \|x\|_k = \sum_{n \in \mathbb{Z}} |x_n| |n|^k < \infty \forall k \in \mathbb{Z}_{\geq 0} \right\}.$$

The standard topology on  $s(\mathbb{Z})$  is determined by the family  $\{\|\cdot\|_k : k \in \mathbb{Z}_{\geq 0}\}$  of norms. Topologize the space  $C^\infty(S^1)$  (by analogy with  $C^\infty[a, b]$ ), and construct a topological isomorphism  $C^\infty(S^1) \cong s(\mathbb{Z})$ . (*Hint:* the isomorphism takes each  $f \in C^\infty(S^1)$  to the sequence of its Fourier coefficients.)

**3.13-B.** Let  $(X, \mu)$  be a finite measure space, and let  $L^0(X, \mu)$  denote the space of equivalence classes of measurable functions on  $X$  (two functions are equivalent if they are equal  $\mu$ -almost everywhere). For each  $f, g \in L^0(X, \mu)$  we let

$$\rho(f, g) = \int_X \min\{|f - g|, 1\} d\mu.$$

Prove that

- (a)  $\rho$  is a metric making  $L^0(X, \mu)$  into a topological vector space;
- (b) a sequence of measurable functions converges in  $L^0(X, \mu)$  iff it converges in measure;
- (c)  $(L^0[0, 1])^* = 0$ .

**3.14.** Let  $\langle X, Y \rangle$  be a dual pair of vector spaces. Show that

- (a)  $\dim X < \infty \iff \dim Y < \infty \iff$  the weak topology  $\sigma(X, Y)$  is normable;
- (b) the weak topology  $\sigma(X, Y)$  is metrizable  $\iff$  the dimension of  $Y$  is at most countable;
- (c) the weak topology on an infinite-dimensional normed space and the weak\* topology on the dual of an infinite-dimensional Banach space are not metrizable.

**3.15.** Describe all continuous linear functionals on  $\mathbb{K}^X$  (where  $X$  is a set), and show that the weak topology on  $\mathbb{K}^X$  is identical to the original topology.

**3.16.** Let  $e_n = (0, \dots, 0, 1, 0, \dots)$ , where 1 is in the  $n$ th slot. Does  $(e_n)$  converge weakly in  $c_0$  and in  $\ell^p$  ( $1 \leq p < \infty$ )?

**3.17.** Give an example of a discontinuous linear operator  $T$  between Hausdorff locally convex spaces  $X$  and  $Y$  such that  $T$  is continuous w.r.t. the weak topologies on  $X$  and  $Y$ .

**3.18. (a)** Give an example of a Banach space  $X$  and a norm closed vector subspace  $Y \subset X^*$  that is not weakly\* closed.

**(b)** Show that, if  $X$  is nonreflexive, then  $X^*$  contains a subspace  $Y$  satisfying (a).

**3.19-B.** Show that each weakly convergent sequence in  $\ell^1$  is norm convergent.